Diving Emergency Action plan 3rd Ed.

Rocket Frog Divers



Let's go make some bubbles

Luis Carlos Huertas Gabert

DAN Instructor #85300

Specialty PADI Instructor #426048

May 2024



OWSI Luis Carlos Huertas

Introduction:

In an area of high practice of recreational diving like playas del coco, located in the north pacific coast of Costa Rica. There are multiples diving centers which give the service of training and recreational diving experience. The probability of an emergency event is directly proportional to the high number of divers in the area, especially in high tourist season which covers the months of December thru June.

Nowadays with the experience of SAR-COV-2 and, airborne pathogens (Virus & bacteria) safety in this regard its important for dive centers with high volume of customers to apply protocols and procedures regarding social distancing, air purity controls, and disinfection of gear in order to keep security and trust of costumer divers based on CDC (Center of Disease Control) and DAN (Divers Alert Network) international guidelines. In this paper we establish a guideline to follow by Boat Staff, Instructors, Dive-masters and DM interns of Rocket Frog Divers.

On the other hand, a diving emergency situation could occur independently if it's a beginner or an experienced diver, if its a newly open Dive Center or an experienced one. Taking this into consideration the application of the Haddon's Matrix model (1) and event can happen if an equilibrium is reached between the variables of the model at any given moment.

It is important for training divers, instructors and general diving personnel to familiarize with the matrix in order to predict and event and avoid it. Another important thing is that a diving center recognizes its limitations, capability of the personnel and stimulate continued training in regards to recognition and providing care of the most common diving emergencies that con turn a recreational diver into a victim.

| | | | Team | Environment |
|---------|--------|---|--|--|
| | | | /Equipment | |
| Phase 1 | BEFORE | -Age -Chronic Medical pathology -Medications or recreational drugs -Training level -Extreme behavior -on edge attitude -Anxiety/stress -temerity | -Trained first response personnel -Full first response equipment - Equipment in optimal conditions -Knowledge on equipment -Crisis manageme nt | -Currents, rain, winds -Visibility -Crisis managemen t of personnel -Fully equipped vessel -Day/night -Available personnel |
| Phase 2 | DURING | -Stress control -Physical condition -Knowledge and reliability on diving equipment -Medical pathology | -Application of action plan -Activation of EMS -Appropriate use of equipment -Most trained personnel managing crisis -Keep calm | -Distance from the boat -Surface, Deep event -Actions of personn el - Communication s |
| Phase 3 | AFTER | -Reanimation effective -Extension of lesions | -Application of guides and protocols -response of the patient | -EMS availability -Distance to adequate health care facility - Availability of resources |

Haddon's Matrix applied to diving (modified by the author)



Even thou this matrix was designed to vehicle accidents its modified version helps the trained personnel to anticipate an event and know how to consider the variables involved in an emergency for them to consider them as RED FLAGS. The more common cause of an incident is poor individual judgment, so this variable is independent of the matrix an involves the diver itself or the actions of one or several parts of the personnel involved in the emergency (here is stated that poor judgment can come from three sides, owner (operator) victim and rescuer) As a rule, the suspicion of a potential risk (red flag) can be assessed and measures can be taken in order to avoid a crisis.

Management of an emergency situation:

An emergency crisis is unexpected even thou there are some variables that can give an index of suspicion. When an emergency occurs, the first rule is: KEEP CALM followed by DO NOT expose yourself to a risky situation (Asses and secure the scene) (2) Keep in mind always be able to rescue yourself first and do not try skills and techniques beyond your capability and training.

If you don't HAVE training as a Rescue Diver, Diving first aid for professional divers (DFA pro) or EMS, do not try to manage the situation, but be ready to assist in the order of your personal training and physical capabilities. Put yourself at the service of the team leader. You still can help holding equipment, securing the area or activating EMS, among other duties that can arise as the emergency develops. Remember an Emergency Scenario is always dynamic, changing second by second, the provider needs to be ready and trained to respond with this concept in mind. Because each situation is different the basic tenet of rescue is "Do and continue doing what works" always looking for the best to the victim without inducing more harm.

The concept of Duty of Care is an obligation imposed on an individual or organization to provide assistance to someone in an effort to prevent unreasonable loss or harm, the care provided should not exceed one's level of training (3)

When providing care treat the injured person with **respect and empathy**, be sure the person is referred to a medical care facility or someone with higher training.

Diving emergencies can occur on dock, deck and sea and in this last scenario two situations can arise; one can be an event on the sea surface or above surface. Each scenario has its complications and needs of management. The personnel proving care should take into consideration each scenario separately and asses the risks involved. Page 5 of

Boat Briefing Guidelines

(table 1) Its imperative that the crew of any vessel be familiar with the emergency procedures and location of emergency devices on land and on board. Without putting aside great costumer service, good service, politeness and a warm welcome it is of extreme importance the safety considerations on land, cars, pangas, on board, during the travel and on the dive site. Our area has different locations which include a national park that has certain regulations to follow an accomplish.

Table 1

Boat Briefing

Welcome to the (Boat name) this boat has certain features (dry storage, area to rest, cooler location and whats on it, bathroom mechanisms, garbage, working or danger area, way of entering the water, way of coming back to the boat, etc) Our staff today are (Introduce captain, instructors, dive-masters, DMT'S etc) In Rocket frog we have an emergency action plan established, the life jackets on this vessel are located : _____ (if captain gives the order we will deploy them to you) the Fire extiguisher is located: _____ the emergency Emergency O2 kit is: _____ We have coordinations with clinics in: _____ and the way to activate EMS is via telephone 911 and channel 16 on Captains radio. Today we are heading to: ______ Distance: ______ Conditions: ______ Safety measures during travel: ______ Captain can ask us to move to certain areas to keep balance of the boat. Once we reach our dive site each guide will give you a briefing for that area.

Other things to consider:

- a- Nearest Hyperbaric chamber (in our case is San Jose)
- b- Special conditions of the dive (negative entry, no line, drift dive)
- c- Responsibility: Level of training, Dive computer, Dive profile, Buddy teams, DECO time.
- d- The Instructor or DM is not a policemen!!!!
- e- Head count
- f- Lost diver procedure (Search for a minute ascend slowly without safety stop) Abort dive for rest of group.
- g- **Recall procedure for divers underwater**: Continuous metal hits on boat ladder and 3 consecutive engine accelerations (as need it).
- h- Captain whistles for:
 - -Abandon ship: six short whistles and a long one
 - -Fire on board: one long whistle (5 10 seg)
 - -Boat danger: One long whistle (5 seg) and 2 shorts

i- Rules:

- DECO (24 HOURS OUT)
- DO NOT PUSH COMPUTERS
- SAFETY STOP 15-30 FEET, 3-5 MIN
- MINIMUM 400 PSI (30 BAR)
- MAX 60 MIN DIVETIME, PEOPLE WAITING
- DEEPEST DIVE FIRST
- DO NOT DIVE ALONE

Its important that the briefing is conducted by the most experience crew member (exceptions BUT ALWAYS supervised is a DMT) and is a duty of all Dive-guides to perform a dive based on the lower level of training of the costumer group (Example: group of 3 divers, two dive-masters and a OW diver with 5 dives, based on this, the dive should be planned based on the OW diver). NEVER a beginner diver should be pushed to perform a dive profile out of his level.

Reinforce that if at the end of the dive anyone is feeling symptoms (3) of:

- FATIGUE
- NAUSEA
- ITCHING
- TINGLING
- NUMBNESS
- PAIN IN ARM, LEGS

Inform right away to the Instructor or DM to proceed with a checkup by a trained staff member.

Captains, Dive Office (administrator) instructor or DM & DMT duties

Besides the regular dive-shop duties and responsibilities of each staff member on daily routine **EVERY** morning Emergency Oxygen kit should be checked (3) by Dive officer, logged and if used it should be filled, cleaned and things used on the First aid kit should be replenished.

Instructors and DM'S should check Tool kit, nitrox analyzer and spare parts in case anything is missed prior to go on the ocean. On board Instructors and DM'S will asses risks on the dive deck and correct them if possible (If not inform directly to Dive officer or captain) they are responsible for gear distribution, tank safety, weights, and costumers on board. They are the eyes of the captain if anything goes wrong during travel.



The dive staff has to stablish roles in case of any emergency, Example:

-**DM 1**: in charge of costumers and asses the scene, helps moving victim, performs CPR if need it.

-**DM 2** : In charge of First aid and O2 kit and if need it delivers O2 if CPR is performed

-**DM 3**: Makes assessments and recollects information if need it delivers first round (2 min) of chest compressions if CPR is performed.

-**Captain**: Safety officer, Communicates with authorities and helps with CPR if need it.

The more staff is on board more duties can be accomplished and it is important that staff is trained in CPR, First Aid, rescue diver, O2 application or Diving First Aid For Professional Divers (DFA pro)

Once on boat everyday Captains should check that life jackets, flotation devices, ropes, anchor, fire extinguisher, Bilge pumps, radio, compass, GPS, lights, fuel pumps, engine, electrical components for any malfunctions or damage.

Captains are in charge of the vessel, personnel are under his command if an emergency is present captain will decide regarding life jackets, abandon ship, switch vessels, create a working area and safety of the boat.

Emergencies and rescue scenarios are unpredictable, even in the safest of environments and with trained personnel, organization, training and simulations are imperative to reduce risks and consequences, post incident debriefing and attending personnel involved on the event is of vital importance to constantly reevaluate protocols and procedures (3)



The general algorithm to approach an emergency is as follows

The first step to respond to an emergency is (Safety) -Captains as a safety officer is in charge of it - secure the scene and by this is making sure that there are no life treading situations for the personnel involved in the rescue and victim(s) management operations. Applying this to the diagram in the first phase which is keep calm, the rescuer or EMS provider must reassure that he has all the equipment necessary to provide help without risking his own life, the rescuer must assess the most secure method to extract the victim out of the water. Once the patient is on the deck the provider must take into consideration biological risks and control them with barriers against biological fluids like blood, vomit, etc.

Every rescue diver, dive-master and diving instructor must keep a proper physical condition, (2) constant training and analysis of medical and non-medical conditions related to diving. Proper training and up to date in Basic Life Support (BLS) Oxygen Provider skills and or Diving First Aid for Professional Divers (DFA pro) equipment and protocols. And one thing to take into consideration is the ability to



ANTICIPATE and detect a probable red flag that can turn a common diving trip to an emergency situation, In other words -DON'T RISK IT-

Any situation that generates stress or anxiety to a diver can turn into an emergency event, there are some signals that can raise the alarm on a trained crew. (2)

- 1- Signals for help (screams, waving hands, mask on the forehead, whistles)
- 2- An evident overexertion on the surface or below.
- 3- Losing pieces of equipment at surface or at the bottom (mask, regulator, BCD)
- 4- Trying to climb desperately and object or person at surface, in order to have a false sense of reach out.
- 5- A total standstill, without any response to a call or help.

According to the diagram, here the rescuer must keep calm and asses which is the safest method to help this distressed diver avoiding first not to get in the water. There are some methods to apply (2):

- A- Reaching and extensions assist: could be in a secure place at deck that the rescuer can reach the victims gear (tank, BCD) without losing grip or could be with an extension like a pole or bar.
- B- Throwing assist: Could be a throw bag attach to a rope, tethered life ring, buoy or personal flotation device. It is important to consider that the object must be thrown past the victim and pull it slowly for its easy reach.
- C- Watercraft assist: Use a small vessel or any other watercraft available to reach the victim.
- D- Wading Assist: In shallow waters, the rescuer enters the water to chest level and tries to reach the victim, it has certain risks cause the entrance in the water can be dangerous depending on the behavior of the victim.

Again, the most important assessment as personnel giving assistance, is your level of training an experience (2,3) it is not recommended to make risky decisions based on good will. Prior to intent any rescue maneuver, the EMS system must be activated cause the basic maneuvers on the deck are not sufficient to any complication secondary to the diving emergency. Based on this principle, the emergency contact numbers in case of any emergency in the area of Rocket frog divers are as follow (table 2)

Once the event is controlled, analysis and **debriefing** with all the staff involved in the rescue or emergency operation is of vital importance to reinforce changes on actions, prevent further situations and asses pos-traumatic stress on staff and victims (3). Debriefing is made **NOT** to charge any one responsible, but more as a learning method and gives the Dive Owners, Dive Officer, Staff in general a tool to improve and makes changes in operation based on evidence. Any unpredictable situation that involves lives creates a general fear feeling of doing something wrong (specially in non professional medical personnel) the debriefing reinforces protocols, detects lack of training and level of training and assess any anxiety triggers and stress created by an event.

Table 2

| Service | Telephone |
|---|---------------------------------------|
| Coast guard | |
| Central office | 2226-3228 |
| | 2226-3238 |
| Flamingo | 2654-6193 |
| lannigo | 20310133 |
| Police of Guanacaste central | 2665-0616 |
| Police of playas del coco | 2670-0258 |
| Police of Flamingo | 2654-6193 |
| BATS islands | 2679–1151 |
| BATS islands Rangers | 2666-0630 ext. 214 |
| Red cross of sardinal (BLS) | 2697-1141 |
| Red cross of Cartagena (BLS) | 2542-5004 |
| Red cross of Santa Cruz (BLS-ACLS) | 2680-0330 |
| Red cross of Liberia (BSL- ACLS) | 2666 0994 |
| Fire department of filadelfia (BLS) | 2688-8733 |
| National Hospital of Liberia | 2690-2300 |
| National Hospital of Nicoya | 2685-5066 |
| National Hospital of Puntarenas | 2630-8000 |
| National central hospital Mexico (san | 2242-6700 |
| jose) | |
| San Rafael Arcangel hospital liberia | 2690-5500 |
| (private) | |
| DAN emergency hot-line | +19196849111 |
| National Bureou of Investigation (OIJ) | 2666-5656 |
| Hiperbaric Chamber Clinica Catolica San | 2246-3000 |
| Jose | |
| Hiperbaric Chamber Hospital CIMA, San | 88306168 |
| Jose | |
| National Aeromedical Service of Police | 2586-4000 / airport of Liberia: 2668- |
| | 1119 |
| Radio Channel for boats emergencies | 16 |
| Badio Channel HF | 8300 MHz |
| Dr Venegas ambulance service -Air | 87771759 |
| Transport- | 0,,,11,35 |
| (Flamingo) GP/BLS/ACLS | |
| National Emergency Services System | 9-1-1 |
| Dr Gerald Smith (EMS physician) | 83465033 |
| SAAT Ambulance Service / air transport | 88289468 |
| (BLS/ACLS) | |
| Dr Mauricio Saldarriaga (EMS physician) | 88356096 |
| Dr Luis Carlos Huertas (EMS physician) | 87189111 |

Emergency Telephone Numbers in the area of Rocket Frog Divers

BLS: Basic Life Support, ACLS: Advanced Cardiac Life Support, EMS: Emergency Medical Services, GP: General Practitioner

As a general rule of information to give to the local authorities, the data must be objective, not suppositions or personal opinions in that regard (2). The general info. Should be given as follows (table 3):

Table 3

Information to authorities

| Hello, my name is:I'm trained in BLS/ACLS (if needed) we are or | n the boat named <u>:</u> | | | | |
|---|---------------------------------|--|--|--|--|
| Number of registry: | I need to report an | | | | |
| emergency related to diving, our emergency condition is: | We are | | | | |
| located: Lat, N, Long W. We are trying to arrive t | to <u>:</u> ETA <u>:</u> At the | | | | |
| beach we need <u>(ACLS unit, oxygen, medical personnel, paramedic, cardiac monitor,</u> | | | | | |
| snake antivenom, transport to hyperbaric chamber) Besides the em | ergency condition of | | | | |
| the patient he has a known medical history of: | <u>.</u> And takes the | | | | |
| following medications: Or any additional information the authorit | zes require. | | | | |

The decision of aeromedical transport is an alternative that is not competitive to the rescue diver, dive-master or instructor. It lays in the physician in charge or that receives the case and manages the patient. It should be based on medical condition, stage of lesions, and weather conditions. In Costa Rica, there are air routes can be taken at 1000 feet is there is visual conditions for flying. An aircraft can fly at 1000 feet in a non-populated zone, and in a populated zone can fly at 1000 feet above the highest point (4). Night flying is not recommended at all (instrumental flying) If ground transport is going to be performed, the ambulance driver must know that ascending from sea level should be at a slow steady pace cause San Jose is at 4921 feet from sea level.

Types of rescue scenarios (2)

1- Tired diver rescue:

A tired diver is a situation that can occur at any moment, and in this scenario a tired diver can turn into a panic diver at any moment; the trigger is unknown. Thou tired divers are usually beginners (he has too many information to process the first times he enters the water) and the overexertion begins at the moment the touch water. Other causes can be underlying medical problems (Arrytmias, Congestive heart failure, medications, diabetics, obesity)

The approach of rescuer must always be defensive. The steps are as follow:

- A- Approach with caution, asses his response and level of calmness
- B- Give direct instructions as to inflate the BCD to acquire proper buoyancy, reinforce to stay calm as the rescuer is approaching.
- C- If the condition of the diver allows, approach and stablish contact thru the back, always speaking to the diver in a calm matter. Secure the divers buoyancy inflating the BCD
- D- If the diver doesn't panic, help him rest, and explain that he is going to be towed to the boat if he wants it; if he can swim the rescuer will escort him to the boat.

Assure the rescuer buoyancy by inflating the BCD.

As said, tired diver scenario can turn into panic diver, specially as evidence shows if the tired diver occurs at surface, its usually described that the victim was not getting enough air, it is decribed as a sensation of breathing through a straw.

If tired diver occurs underwater TIME is the key, rescuer must try to calm the diver as fast and steady as possible to be successfull.

2- Panic diver rescue:

In order to perform a proper panic diver rescue, the approach must be cautious and slow, always assessing the diver metal state in the same matter as a tired diver. The panic diver doesn't obey orders, tries to take out his equipment and is a dangerous victim cause this diver will try to hold on to whatever is near in order to keep buoyancy. The rescuer must be in control and secure when he approaches this scenario. The approach must be thru the back or from under the surface, grabbing his wrist and turning him, then holding the tank and securing buoyancy inflating the BCD, and then assuring buoyancy for itself. If it's impossible to approach to the panic diver, maintain a considerable safe distance an let the diver get tired or turn unconscious and make a cautiously approach. Always talk to



the diver, assuring that the rescuer is there, and trying to calm him down. When on hold to the diver, toad him to the boat in the same matter as the tired diver.

If panic diver occurs underwater, DO NOT TRY TO KEEP regulator in mouth, if he can't breathe at the point of approach he will continue with difficulty breathing. TIME is definite concern, panicked diver tissues will continue to consume oxygen, even when he is not breathing. During the ascent, as oxygen concentrations in lung decrease and gas expands the diver is at risk of losing consciousness (Hypoxia of ascent or "shallow water blackout") risky situation of arterial gas embolism from lung overexpansion.

3- Unresponsive diver:

In this scenario the approach should be cautious, when in contact asses the level of consciousness of the diver, if his head is below the surface it should be turned up and secure the airway to assess airway permeability and breathing. If the victim is unresponsive, call for help with signals, whistle or yelling.

If the rescuer can transport the victim in less than five minutes to the deck, the rescues should give two rescue breaths every five seconds assessing permeability and swimming the victim to the boat. If the boat or secure area is more than five minutes away give rescue breaths during a minute and if he continues unresponsive swim as fast as possible to a secure area. Once on deck start BLS or CPR and emergency response protocol.

If diver is unresponsive underwater TIME IS THE KEY, if he has regulator on mouth hold and ascent on a pace not fast than 18mts / 60 feet per minute. If he doesn't have regulator in mouth DO NOT TRY to put it back.

If diver is unconscious he may relax the pharynx and allow passive exhalation, this will allow air to vent out the lungs without over-pressurization and barotrauma. The opposite may occur an may experience laryngospam (a reflex spam that occurs when water enters throat) this can cause a partial or complete closure of the airway an during ascent this reflex presents the risk of arterial gas embolism.

4- Lost diver scenario

When a lost diver scenario appears during a dive, it is important to assess the situation as quickly as possible, the most probable condition will be death. So, every second counts. This scenario is resource dependent; the steps are as follow:

- A- Alert EMS
- B- Detect where was the last area the diver was seen
- C- Delegate watchers on a high point to detect bubbles on the surface.
- D- Try to find out if the diver came out first and left the diving site on his own (search for his equipment)

- E- If divers are available, send them to the last area where he was seen
- F- Mark with a buoy the last area he was seen
- G- Send snorquellers to scan the area

A lost diver in the first minute without air cardiac complications appear, like arrythmias and syncope. At this stage the level of consciousness of the diver is lost. In between five to ten minutes irreversible brain lesion appear and death is a certain situation. In this scenario the quickness of the team is a priority, always taking into consideration DO NOT risk unqualified personnel and don't take risks for a dead body.

5- Vessel in Danger

This scenario is relative to risk of collision, fire, sinking or other. It prepares the crew and passengers to an emergency situation. Captain raise the alarm with **one long whistle and two shorts**. An informs to main office and authorities of the situation. Crew stays at ease to captain commands and prepares emergency stations as need it (02 Kit, Life jackets, fire extinguisher, flotation devices, etc) and assures the costumers are in a safe area or position.

6- Abandon ship scenario

This type of emergency is catastrophic, the boat is lost and is going to sink. But injuries can be reduced if personnel is well trained. In this scenario the captain is the ONLY makes this decision, and informs the main office and crew of what is happening, coordinates another rescue vessel and speaks with authorities. Cause the crew has assigned roles prior of departing let's consider this scenario with a 3 persons crew boat:

-**Captain**: Gives the order to abandon ship **(six loud whistles and a long one)** informs main office to coordinate rescue and alerts authorities via 911 or channel 16.

-**DM 1**: Deploys life jackets, makes sure all costumers put their life jackets and are on safety positions to go into the water. Keeps closed eye on water level and risks on the conditions the boat is. Once on the water keeps together all people, holding and floating while rescue arrives. Waits for orders of the captain.

-**DM2**: Helps DM1 deploying life jackets, organizes all costumers to abandon ship. Keeps a head count and informs DM1 and Captain, assists DM 1 in keeping people together on surface and floating. Waits for orders of the captain.

Its imperative all staff members perform drills and resfresh the concepts of the role assignment and be up to date on administrative changes.

7- Fire

Any person on the vessel, costumer or crew can detect a fire, it is of vital importance for everybody on board to know where fire extinguisher is. The alarm is raised with a **constant whistle at least 10 seg.** Captain changes course in direction of wind, disconnects electrical devices and shuts engine. DM's are under command of captain and make sure costumers are safe and protected, while fire is being controlled. Of important matter is location of O2 kit cause if makes contact with fire is of extreme risk.

All scenarios can mix and can evolve from one to another, **MUST REMEMBER**: An Emergency is Dynamic, so it is important to consider training, simulation and constant interaction within the crew to overcome any unpredictable situation.

Diving Emergencies

Start of the Dive:

When a diver enters the water, his body responds to the change in temperature, including a reduction of peripheral blood flow to maintain core temperature. The reaction may also include a gasp for breath, which could, in extremely cold water, escalate into hyperventilation, overexertion and drowning. Immersion can cause pulmonary edema, arrhythmias leading to syncope or death. Death is rare from the initial immersion, but if an emergency is developed at the beginning of a dive, a reaction to initial immersion should be kept in mind (5) In this regard it's important for the rescue diver, dive-master or instructor to assess the divers face and reactions at surface, asking a few simple questions could asses this condition. MUST CONSIDER THAT a foreign or local diver may not want to express his discomfort or peer pressure risking his dive day.

General Environmental Considerations

1- <u>Hypothermia, Hyperthermia, IPE & Dehydration:</u>

Although divers take steps to keep warm while diving, heat loss will occur. Divers on vacation often make multiple dives per day and subject themselves to prolonged cooling over the course of the day. Further heat loss (and gain) can occur as divers are exposed to the sun or wind between dives (5)

Diving also causes "immersion diuresis," which can lead to dehydration and associated issues, including syncope and arrhythmias (5) Dehydration is exacerbated by other factors, including carrying equipment, exposure to sun, breathing dry, compressed air, airplane travel and taking diuretic medications as chronic treatment.

Hypothermia, hyperthermia and dehydration should be treated in accordance with your local protocols or recommendations of primary physician or EMT. Keep in mind that certain signs and symptoms of these conditions, such as nausea, are also associated with other diving injuries. Dehydration should be treated with electrolyte oral solution (bottled or powder mix) not with plain water (risk of Hyponatremia – Low sodium-) (3)

Another condition **Immersion pulmonary edema** (IPE) a term used by many as an umbrella term for scuba divers' pulmonary edema (SDPE) and swimming-induced pulmonary edema (SIPE). Both of these conditions share similar presenting symptoms, diagnostics, and treatment (6,7) Case studies from the 1980s to the present time highlight differences in pathophysiology (cause and mechanism) and certain predisposing factors. Both conditions are likely underdiagnosed since postmortem exams with pulmonary edema exhibit nonspecific findings of heavy edematous (swelling) lungs and frothy sputum (pink, bubbled flem) in the airways (5) These same findings can be found following drowning and prolonged resuscitation (CPR) efforts.

There are numerous predisposing factors to immersion pulmonary edema, for example, overhydration, cold water, negative inspiratory pressure, asthma, diabetes, drugs (such as beta-blockers), underlying cardiac problems, physical exertion, previous episodes of IPE, and systemic and pulmonary hypertension. (5) Cold water is believed to be a major contributing factor to the underlying pathophysiology. However, there have been numerous case reports and studies with individuals diving in tropical waters of 30 to 35 C. The majority of the incidence of IPE in warm water was diving-related, but it has been seen with extreme exertion and overhydration in swimmers.(5,6,7)



Barotrauma

In much the same way you may feel ear pain during take-off or landing during a flight due to pressure changes, the changes in pressure when diving can cause injury to ears, sinuses and other air spaces. Although decompression illness and other emergencies are dramatic, ear injuries are much more common.

1- Ears & Sinuses

Divers are trained to equalize the pressure in their ears to ensure that air enters the middle ear through the eustachian tubes (8) Pressure may be felt quickly, and a 10- to 12-inch descent can be sufficient to cause discomfort and pain. Barotitis media (middle ear barotrauma) occurs when pressure in the middle ear is lower than the surrounding pressure (which may occur during descent), forcing the eardrum inward and causing pain and bleeding. A pressure differential can also cause the eardrum to tear or rupture, which results in pain and bleeding. Water entering the ear may cause vertigo, which is very dangerous when it occurs at depth (5) The membranes in the sinuses can also be damaged by pressure. Note that sinus barotrauma and related injuries can occur both during descent and ascent.

When ascending, the mechanism of injury is reversed as air expands called reverse block.

During ascent and descent, the rescue diver must observe the divers, asses if there is abnormal behavior, and assist any distress diver. If the afflicted diver can't correct problem abort the dive and asses.

An injury to the ears will manifest as a feeling of fullness, pressure or pain. There may also be a hearing loss or complaints of a "squeaking sound." Blood or other fluid discharge from the ear may be evident. A sinus injury can also cause fullness, pressure or pain (including pain in the upper teeth) with blood or other discharge from the nose.

Divers with suspected barotrauma should not return to the water(3,9) They need to be assessed by a physician for proper treatment, which may include antibiotics, antihistamines and in severe cases surgery.

2- Effects on the Lungs

The effects of the change in air volume and the other gas laws form the basis of decompression illness (DCI), which includes arterial gas embolism (AGE), and decompression sickness (DCS), commonly known as "the bends." (8)

If a diver takes a breath of air and starts to ascend while holding his breath (or with air otherwise trapped in the lungs), the air expands as the ambient water pressure decreases, much like air expanding in a balloon. This can damage the alveoli and bronchial passages.

Expanding air may rupture the lung and cause a pneumothorax. Mediastinal and subcutaneous emphysema can result from air that escapes from the lung. This condition is known as Lung Overexpansion injury (7).

AGE occurs when bubbles are forced into the bloodstream through alveoli, resulting in blockages when the bubbles reach smaller blood vessels. A cerebral arterial gas embolism (CAGE) results if the air embolizes in the brain (10)

AGE will usually occur either during a dive or within 15 minutes after a diver surfaces (8, 11) Symptoms include chest, torso or back pain, malaise (feeling unwell) nausea, coughing and shortness of breath, bloody or frothy sputum, headaches and/or dizziness, rash visual disturbances (including partial or full blindness), numbness and/or tingling (in abnormal areas of body – nose tip, elbows, belly, knee) weakness or paralysis and loss of sensation in the body, and syncope (8).

Other atypical sings and symptoms (9, 10) are blochy skins -Livedo reticularis-, confusion, convulsions, difficulty passing urine, lack of coordination, itch, loss of baldder / bowel control, personality change, tinnitus (ringing in ears) speech disturbances.

Neurological symptoms are present aprox. 55% cases. Mixture of symptoms with neurological affectation occurs in 80% of cases (8,9).

ANY SIGN OR SYMPTOM THAT OCCURS WITHIN 24 - 48 HR OF DIVING should be considered diving related until proven otherwise. Especially if the depth-time exposure has approached or exceeded accepted procedures (10,11,12).

a. DCS: "The Bends"

As a diver descends, the increased ambient pressure forces inspired nitrogen into tissue. (11) When a diver ascends the pressure is reduced and nitrogen is released from tissue into the bloodstream. If a diver ascends too quickly, has been down too long or otherwise has excessive amounts of nitrogen in his body, bubbles may form in the bloodstream or remain trapped in tissue. (12)



Since bubble formation can occur in any tissue, there are many forms of DCS. Pain in recreational divers is more common in the arms, whereas in saturation divers the pains is in the knees. Either may occur in the limbs or joints. Central nervous system DCS can cause paralysis and loss of sensation. Cerebral DCS may produce headaches, visual disturbances, paralysis, unconsciousness and altered mental status. "Skin bends," or cutaneous DCS, present with itching, burning or mottling of the skin (10,11).

The onset time of DCS varies within 24 hours of diving. In a recent report, half of all cases of DCS had symptom onset times of an hour and a half or less after a diver surfaced. Any symptoms that occur after 24 hours are probably unrelated to diving. In rare cases, divers exposed to altitude may exhibit initial symptom onset after 24 hours (10).

DCI initial treatment on deck

The basic treatment for suspected cases of any DCI, including AGE and DCS, are essentially the same, with the general algorithm of emergencies the assessment of ABCs is a priority. The importance of administering 100% oxygen cannot be over-emphasized in suspected cases of DCI or other diving NO CLEAR etiology incidents (10) (This is why all staff members should be familiarized with the O2 kit and the way it fully works) The reasons to apply 02 include the following:

Reasons and considerations using oxygen therapy (10)

| 1- | Breathing 100% 0xigen is the first aid for diving incidents |
|----|--|
| 2- | High concentration oxygen flushes out dissolved nitrogen and nitrogen |
| | bubbles and supplies oxygen to hypoxic tissues |
| | a- This gives the advantage that if incident is not diving related clarifies |
| | the clinical spectrum of the patient ruling out nitrogen bubble formation |
| 3- | Oxygen breathing should begin as soon as possible until a medical |
| | professional request it to |
| | be ceased |
| 4- | It is important not to stop oxygen breathing to soon. Even if initial |
| | symptoms have |
| | disappeared, they can re-appear later and can worsen substantially |
| 5- | If a demand valve with mouth piece is used on conscious diver, their nose |
| | must be blocked |
| 6- | All dive operators should have suitable oxygen equipment for divers and |
| | staff trained and |
| | competent on its use |



In addition to providing respiratory support as it does in any emergency, oxygen promotes the elimination of inert gas bubbles in the tissues. Dive boats often have oxygen on hand, and someone may have begun administering oxygen prior to EMS's arrival. DCI can be life-threatening, and any suspected DCI requires immediate transport to a medical facility where a complete diagnosis and treatment, **WITH AVAILABILITY OF** hyperbaric chamber therapy. Note that changes in altitude can exacerbate a case of DCI, so if the patient needs to be transported by helicopter, it should be at the lowest safe altitude possible. The current recommendation is 1000 feet' or lower (10, 4)

A physical exam, primary and secondary evaluation, including a thorough neurological examination, should be performed and changes should be monitored carefully. NEVER ATTEMPT "recompression" by placing a diver back into the water(8,9).

| Finger nose movement | a- eyes open b- eyes closed |
|---------------------------------|--------------------------------|
| Ability to stand | a- eyes open b- eyes closed |
| Walking heel-toe short distance | a- front b- back |
| Ability to walk free | Walk on deck |

Neurological test that can be fast performed on deck (3,10)

Always perform test supporting patient, careful with fainting and falling

Part of neurological testing is responsiveness of the victim (3)

- 1- alert
- 2- Responds to verbal stimuli
- 3- Responds to painful stimuli
- 4-Unresponsive.

Other important concept is if the victim is Oriented in person, time and place and how is his/her speech, abstract reasoning, calculations and short term memory (3). Neurological assessment is a skill trained in courses like Diving First Aid for Professional Dives (DFA pro) and on Primary and Secondary assessment courses. Of vital importance is to check for Fascial drop, Arm or extremity weakness, speech difficulty or sudden severe headache and keep a record of the beginning and duration of this symptoms not forgetting to call EMS (911). Neurological afflictions secondary to Scuba dive are very subtle (3,9,12) and usually serious so when suspecting one its important to take into consideration medical history of the victim like:

- 1- Signs/symptoms
- 2- Allergies
- 3-Medications
- 4-Pertinent medical history
- 5- Last oral intake
- 6- Events leading to current situation.

3- <u>Shock</u>

To the BLS provider shock could be confusing, it's a myriad of signs and symptoms that are the Lobby to organ failure (9,12). Usually when data of shock appears, the physiological celular clock is suffering and the battle is 50% lost. The most common scenario for shock is bleeding, in which the patient enters to a state of hemorrhagic shock due to blood loss. The signs and symptoms of shock are as follows on table 4.

Table 4

| Signs | Symptoms | | | |
|--|---|--|--|--|
| Pale skin | Cold | | | |
| Cold levels in extremities | Cold sweating / clammy skin | | | |
| Anuria (don't urinate) | Dry mouth | | | |
| Capillary filling more than 3 seg | Thirst | | | |
| Radial pulse absent | Taquicardia / Palpitations / Arrythmia | | | |
| Cyanosis (Purple color in fingernails) | Restlessness / irritability / Incoherence | | | |
| Pulse 02 below 94% | Shivering | | | |
| Taquipnea (rapid breath) | Dyspnea (sensation of shortness of | | | |
| | breath) | | | |
| Dazed look | Nausea / Vomiting | | | |
| Syncope | Loss of consciousness | | | |

Signs and symptoms of shock

As BLS the most important maneuvers a rescuer can provide is control of bleeding, give oxygen in high percentage and control hypothermia. There's a triad in which the patient in shock should never get into that is called the triad of dead (Hypothermia, Acidosis and coagulopathy) So in a boat, ending a dive, the patient should be removed of all wet clothing, maintained warm with all the resources possible until it reaches ACLS care. If there's an ACLS provider on the boat, the most important thing to do (besides the latter) is apply IV fluids, raise systolic blood pressure and preserve kidney function in other to eliminate the metabolites that are produced in the shock state.

Breathing Dangers

1- Nitrogen Narcosis

Inert gases under pressure have an intoxicating effect. Nitrogen under pressure can cause nitrogen narcosis, which impairs the ability to think and reason. Divers can usually stop nitrogen narcosis by ascending until symptoms resolve. The danger of nitrogen narcosis lies in impaired judgment, causing divers to lose track of time, remaining air and depth limits and resulting in agitation, panic or confusion. This can lead to various problems, including drowning when they run out of air, or increasing the chances for serious injury or illness by ascending too rapidly. (12) Divers can recognize this state by watching the diver with erratic behavior.

2- <u>Oxygen Toxicity</u>

Divers run little risk of oxygen toxicity when they stay within recreational depth limits of 100 feet and use dive regular air, but it becomes a concern when they use mixed gases like enriched air nitrox (11,13) When using EAN-32 its important to understand dive profile, ceiling and DECO time concepts. There are two types of oxygen toxicity. The first is central nervous system (CNS) oxygen toxicity. Signs and symptoms of CNS oxygen toxicity include convulsion, visual disturbance, tinnitus, nausea, tingling, twitching, muscle spasms (especially along lips and mouth), irritability, dizziness and/or dyspnea. CNS toxicity is the more dangerous form of oxygen toxicity due to the risk of having convulsions under water (3,8,11).

The second type is pulmonary toxicity, which can affect the lungs or other parts of the body. Symptoms include chest pain and discomfort, coughing and fluid in the lungs.

3- Carbon Monoxide Poisoning

Carbon monoxide poisoning can occur when the air in a scuba tank has been contaminated by fumes from an improperly maintained air compressor. This is why is important to assess the odor of the air prior to assemble the scuba equipment on the boat (2,3,11) Carbon monoxide poisoning often produces no symptoms before



the diver loses consciousness, although there may be symptoms including headache, dizziness, nausea or altered mental status. There may also be excessive red or blue coloring of the lips, nail beds or skin. Patients with suspected carbon monoxide poisoning should be given oxygen.

It's important for the rescue diver, dive-master and instructor to know that there are going to be questions that a trained physician or investigator could ask On Scene

As in any other emergency, fact-gathering is important. Divers are trained to dive in a way that reduces the chance of DCI or other injury, but they may still experience problems even if they follow all the rules. In addition to medical history and other information you normally gather, there is information particular to diving that treating physicians will find useful.

Questions that could be asked and should receive an answer include:

- a- "How long did it take the diver to ascend to the surface, and did he stop along the way?" A slower controlled ascent coupled with one or more stops helps the body off gas before bubbles form and also helps prevent barotraumas from air expansion.
- b- "How deep and long was the dive?" There are recommendations on how long a diver may remain at various depths in order to minimize the risk of bubble formation.
- c- "How long was it before the diver showed signs or symptoms?" DCI (including different forms of DCS) will present at various times based on severity and the type of DCI.
- d- "What type of breathing gas was the diver using?" The gas mixture can help determine what illness or injury occurred. Divers are normally trained to dive with a buddy, who may be able to provide information.

As a legal concern, IT IS IMPORTANT TO don't interfere with the equipment the diver was using other than to close the valves on the tank and note the number of turns it took to close the valve. (16) Another important consideration is his diving computer. IT SHOULD NOT BE TOUCHED. There may be regional differences regarding what actions should be taken, including whether you should close the valves on a tank, so follow local procedures and defer to the judgment of law enforcement or other investigators on scene.



Disinfection and Gear Cleaning

Even thou Scuba Diving gear with proper cleaning and maintenance is a low risk transmitter of airborne diseases, Rocket Frog divers has a protocol in regards to cleaning disinfecting and drying 2nd stages and Wetsuits in an area of the shop specific for this purpose. Individual prevention is based in avoiding (16)

- 1- Direct contact with respiratory fluids of an infected symptomatic or asymptomatic patient.
- 2- Disinfecting hands and portables with soap and water.
- 3- Avoiding shaking hands, keeping prudent space between people.
- 4- Avoid touching eyes, mouth and nose without washed hands
- 5- Being careful and responsible while coughing and having flu like symptoms

There is no direct evidence of the presence of virus and bacteria in dive gear, hoses, second stages or primary stages. By this DAN (divers alerts network) gives guidelines of cleaning gear with proven disinfectants (which can be found on CDC and EPA web page) (15, 16) that do not damage neoprene, plastic, metal. Like sodium Hipochlorite (Clorox) or quaternary ammonium, both diluted in water (16).

- 1- DAN recommends submerge gear for 5 min or more on water with Sodium Hipochlorite.
- 2- Clean and wash gear on a ventilated area
- 3- Mix active ingredient with cold water only (hot water damages active principle)
- 4- Use gloves, eye and mouth protection when mixing water with sodium hypochlorite or quaternary ammonium.
 - a. 5% : 100 cc of Sodium Hypochlorite in 5 liters(5000cc) of water

b. 10%: 50 cc of Sodium Hypochlorite in 4.9 liters (4900cc) of water (recommended) 5- When finishing cleaning gear with active principle wash thoroughly with clear fresh water.

With air purity and quality of compressors, Rocket Frog has a maintenance team specific for this tasks, membrane and Nitrox (EAN) quality is checked every day by rigid supervision and CO levels are measured multiple times during the filling process. Tank inspection and Hydro test are performed by qualified personnel and if need it third party inspection is performed.



As a rule staff must recognize a costumer with flu like symptoms and ask general questions to report to Course director, shop administrator or CEO:

- 1- Presence of fever
- 2- Cough, lost of taste or smell on last days
- 3- Muscle pain, diarrhea, dry throat
- 4- Using any over the counter medications

Marine Life Injuries

Marine life injuries are frequent, as a rule bad judgment dictates touching marine life and dab training brings bad trim and buoyancy. Considering how many organisms in the marine environment can cause harm to humans. Divers can prevent marine life injuries by adhering to the general diving rule, **"no touching, teasing or taking."** (3, 14) Besides encouraging using gloves, wetsuit, and rash guard is a good preventable measure. Sometimes, however, contact with harmful flora or fauna underwater is either unexpected or unavoidable. In these cases, it's important to know how to react. In minor cases, delivering effective first aid on deck provides pain relief. In more serious cases it can vastly improve a victim's chances of survival. Here we'll take a look at four potential marine life injuries and the first-aid procedures required to treat each one (3, 12).

1- Jellyfish stings

Of the myriad jellyfish species, some are sting-less and some can be fatal. When stung by a jellyfish, the seriousness of the situation depends on several factors. These include the species, the anatomical location of the sting, the size of the affected area, and whether or not the victim is allergic to the animal's toxins. Many jellyfish stings do not require professional medical attention. But if the victim experiences severe symptoms, including chest pains and difficulty breathing, contact emergency medical services immediately.

If the victim is particularly old or young, or the sting covers a large area, call emergency services immediately (3, 11,12). In general, however, first aid for all jellyfish stings follows the same guidelines. The victim should be treated out of the water, and kept as still as possible. This prevents more toxins from being released and spreading around the body (12).

If there are any visible tentacles on the victim's body, remove them using tweezers, forceps, or gloves to avoid further stings. After you have removed the

tentacles, clear any remaining stinging cells, or nematocysts, by applying shaving cream. Use a razor or credit card to carefully scrape them from the skin. If you do not have these tools on hand, use salt water to flush the affected area. Never use fresh water, as it can cause unfired nematocysts to release their poison (2,3,14).

Contrary to popular urban myth, you should avoid urine for the same reason, as well as alcohol. Some sources (including DAN) recommend using household vinegar for sting relief. Others, such as the British National Health Service, advise not to for fear that the vinegar may also trigger remaining nematocysts. Instead, they recommend either applying an ice pack to the affected area or soaking it in hot water. DO NOT scrub the area, just irrigate. This offers the victim pain relief after all tentacles and stinging cells have been removed (3, 14).

2- Lionfish, scorpionfish and stonefish injuries

All three of these fish species are highly venomous. They all have spines on their dorsal, anal and pectoral fins, capable of injecting venom into a diver's skin. Typically, injuries from any one of these fish require emergency care. If you are stung by a stonefish, seek professional medical attention immediately (3, 8). Of the three, the venom of this fish is the most potent and can be fatal. Victims will require anti-venom injections as part of their treatment. First-aid care is the same for both lionfish and scorpionfish injuries.

Once the diver is safely out of the water and in a stable environment, use tweezers to gently remove any spine fragments embedded in the wound. Then, soak the wound in hot water. It should be as hot as the victim can tolerate without burning the skin. These species' venom is protein-based, and begins to break down and deactivate with the application of heat. Soaking the wound in hot water will not only relieve pain but also reduce the venom's effects (3, 14).

If possible, immerse the affected area for at least 30 minutes. If the wound is located somewhere that makes immersion difficult, use hot washcloths. Hot-water treatment may also help in the case of stonefish injuries. Apply hot washcloths or immerse the wound while waiting for medical care to arrive. After soaking, disinfect the wound, apply antiseptic ointment and take over-the-counter pain medication to help alleviate pain. Divers should seek immediate medical attention after exposure to the venom of any of these species, as some complications may take time to manifest. Serious complications include anaphylactic shock as a result of an allergic reaction, shock, infection, tissue death and paralysis. These wounds can take weeks, or even months, to heal (3, 11,12). Under medical supervision you can give antihistamines, paracetamol orally if not contraindicated.

3- Coral cuts and abrasions

These are perhaps the most common diver injuries. Coral scrapes can occur whenever a diver contacts the reef. Coral is often sharp, and those who get cut will find that they take a long time to heal, and often become infected. This happens because a thin film of living organisms covers the coral. They tear away from the main structure on contact and contaminate the victim's wound. To treat coral cuts and abrasions, first stem any significant blood flow using direct pressure. Once the bleeding has stopped, make sure to remove any remaining coral fragments by flushing with clean fresh water. Use antibacterial soap or hydrogen peroxide mixed with water to disinfect the wound. Finally, rinse again with fresh water (14).

After you have cleaned the wound, apply antibiotic cream. Cover with a sterile, non-adhesive dressing. Clean and re-dress the affected area twice a day until the wound has healed. Although victims won't usually require emergency medical care for coral injuries, watch closely for signs of shock or aggravated infection. Treat extreme redness, swelling, excess pus, swollen lymph glands or any other signs of fever or infection immediately.

4- Large animal bites

When it comes to marine life injuries, most people automatically imagine shark

attacks. Truthfully, the likelihood of receiving a shark bite is infinitesimally small. But several marine species can inflict a nasty bite, including moray eels, barracuda and seals. Usually these species will only react aggressively in self-defense, or if they mistake a diver for prey. The severity of such injuries can range from the superficial to the fatal. In the case of more serious bites, first remove the victim from the water (3,12,14).

Contact emergency services immediately, and then focus on providing basic life support until the professionals can take over. Apply pressure to the wound to stem blood flow, remembering to protect yourself wearing protective barriers if possible (latex gloves, goggles). Do not release pressure to change bandages or cloths even if they become saturated. Instead, simply apply more bandages directly on top of the original ones (11). If the bleeding is excessive and you cannot stanch it, apply a tourniquet as close to the injury as possible. If not used properly a tourniquet can cause other complications.

Tourniquets are an absolute last resort, and ideally someone with prior medical training will apply them (14).

If the bite is located on an extremity, raise it above heart level to help reduce blood flow. If the bite broke a bone, use a splint to support and protect the injured limb. If you don't have a purpose- made splint on hand, you should be able to improvise one quite easily. Often, the shock and trauma of a severe bite can cause a victim to have difficulty breathing; in this case, administer oxygen and monitor continued respiration until emergency services arrive. In the event of minor bites,



the victim may not require emergency services. Simply use direct pressure until the bleeding stops, and then rinse gently with fresh water, clean with antiseptic and apply a non-adhesive, sterile dressing. Should any infection occur, or other symptoms arise, make sure to seek medical advice immediately (14).

5- Stingrays

Stingrays are generally docile creatures, but are equipped with a tail that has sharp barbs used for defense, most often on an unsuspecting person who has the misfortune to step on it. The tail inflicts a deep puncture and/or laceration together with contact to the proteins the barbs has on its tissue. The pain is often severe and accompanied by bleeding. Reactions include weakness, fainting and, in more extreme cases, anaphylaxis, paralysis and death. You should first establish the universal algorithm of assessing the ABCs, then rinse the wound with clean water and immerse in hot water (110°F-113°F/43.3°C-45°C) for 30 to 90 minutes. Extract obvious pieces of barbs, then conduct pain management as needed (14) As with any wound, there is a risk of infection from the puncture. Management under medical supervision with antihistamines, paracetamol and antiseptic soap is advised if victim is not allergic.

6- Marine Snake Bite

The Hydrophis platurus (yellow belied sea snake) an its variation found in the Golfo de Nicoya H. platurus xanthos, is a small headed snake of the same family of the Elapidae sp. or Coral snake. The venom is neurotoxic, there are no reports on medical literature of a Hydrophidae attack on humans. There is no antivenom for this snake, it is theorized that the anticoral serum on high doses could be a good therapeutic to this event, thou more evidence is needed. This snake is afraid of humans, not aggressive and usually runs away. But it should not be bother as any other marine life, she could attack on self defense and because her jaw is small, areas like hands, ears, nose and feet are probable targets.

Dive plan to prevent emergency events

Dive plans set out the way diving will be carried out to ensure the participant's health and safety. The dive plan should assist with the implementation of control measures that have been selected during the risk management process.

A single dive plan may be prepared for several dives when the risks of the dives are similar. When the risks of the dive vary (e.g. when there are different tasks, equipment used or significant environmental change), then a new dive plan should be prepared.

Simulation, role assignment, and training is imperative for the effectiveness of the dive plan.

Example dive plan for diving work:

| Dive plan | | | | |
|---|-------------------------------------|------------------------|----------------------|-------------------------------|
| Business legal ABN: name: | | Depart date: | | Return date: |
| Dive supervisor: | | Vessel master if a | | licable: |
| Dive site locations: | | • | | |
| Dive plan prepared by: Date: | | | | |
| All persons briefed by d | ive supervisor | | | |
| Method of carrying out the | task | | | |
| Type of diving | Tasks | | Diving equipment/gas | |
| ै Vessel - per ops manua | ै Manual clea | aning | ို SCU mai | BA air only - per ops nual |
| ီ Shore - per ops manual | ै Harvesting | by hand | ै SSB mai | A air only - per ops nual |
| 2 | ै Inspection | | ໍ້? Pre ແ com | leparture checks |
| ै Scient | | ampling | ै Pr availa | e dive checklist ble |
| ŝ | ð | ້ຳ 🕴 🕈 Other (specify) | | er (specify) |
| Decompression management | | | | |
| ै Computer (specify) | ້ຳ Tables (specify) ້ຳ Flying after | | ng after diving | |
| Other deco factors (specify) | | | | |

| Task and duties (all members of dive team) | | | | | |
|--|-----------|---------|--------|--|--|
| Name | Competent | Medical | Duties | | |
| | | | | | |
| | | | | | |
| | | | | | |
| | | | | | |

| Site specific hazards (in addition to controls in ops manual) | | | | | | |
|---|-------------|-------------|----------|-------------------------------|--|--|
| Hazard | | Risk | | Control measures (site rules) | | |
| Environmental hazards | | | | | | |
| (specify) | | | | | | |
| Equipmenthazards(specify) | | | | | | |
| Task hazard | s (specify) | | | | | |
| ै Other (speci | fy) | | | | | |
| | | | | | | |
| | | | | | | |
| | | | | | | |
| | | | | | | |
| Diving planne | d | | | | | |
| Date range | Location | Depth range | Max time | Task | | |
| | | | | | | |
| | | | | | | |
| | | | | | | |
| | | | | | | |
| | | | | | | |
| Emergency procedures | | | | | | |
| Emergency procedures available on site (as per ops manual) | | | | | | |
| Emergency procedures reviewed and practiced by dive team | | | | | | |
| Last drill date: | | | | | | |

Applying to the plan and getting knowledge of it by all members of the crew reduces the incidence on events, and prevents an emergency.

Page 23 of

Conclusions:

Medicine is an ever-changing scientific area, managements and protocols are changing day by day that's why continuous reading, updating and training is needed. This document should not be considered a training and or training course. In a vast area as diving, which has so much to investigate still, the science of injury and pathology is ever changing. Divers Alert Nertwork annual reports gives insight onto the evidence of diving medicine and its advances. The need for evidence-based studies and reports is of vital importance. As a high-volume dive center, reports, statitics and incidence of cases could make a big difference toward the changing field of medicine and diving.

In regards to emergent infectious diseases people at high risk should follow health services recommendations, control their chronic established diseases and maintain strict hygiene measures. Rocket Frog divers in this regard, keeps rigid protocols with air filters, membrane maintenance and tank inspection.

Finally, because of the high volume of divers Rocket Frog has, the continuous training, simulations, studying and applying the knowledge is imperative. Accidents and events occur proportionally to the volume of activity in a field.

Bibliography:

- [1] Runyan CW. Using the Haddon matrix: introducing the third dimension. Inj Prev. 1998;4(4):302–307.
- [2] PADI, Rescue diver manual, PADI Publishing, 2016
- [3] DAN, Diving First Aid for Professional Divers, Student Handbook, Divers Alert Network, 4th edition, Rev. 3.0, 2023
- [4] Direccion general de aviacion civil Costa Rica. Lineamientos de CO para aeronaves ala fija y ala rotatoria. 2008.
- [5] Douglas JDM. Medical problems of sport diving. Br med J 1985;291:1224 /6.
- [6] Kumar M, Thompson PD. A literature review of immersion pulmonary edema. Phys Sportsmed. 2019 May;47(2):148-151. [PubMed: <u>30403902</u>]
- [7] Castagna O, de Maistre S, Schmid B, Caudal D, Regnard J. Immersion pulmonary oedema in a healthy diver not exposed to cold or strenuous exercise. Diving Hyperb Med. 2018 Mar 31;48(1):40-44. [PMC free article: PMC6467827] [PubMed: 29557101]
- [8] Sykes JJW. The hazards of sport diving. J Roy Coll Physicians Ed 1990;20:318/22.
- [9] Strauss MB, Borer RC, Jr.. Diving medicine: contemporary topics and their controversies. Am J Emer Med 2001;19:232/8.
- [10] Vann RD. et. al. Decompression illness. LANCET mag, 2011: vol 377 (Jan 8): 153-64 [8]Dovenbarger J, Uguccioni D, Report on decompression illness and diving fatalities, Durham, N C, Divers Alert Network. 1999
- [11] Lippmann J. Decompression Illness. J.L pubications.
 2011.Melamed Y, Shupak A, Bitterman H. Medical problems associated with underwater diving. N Engl J Med 1992;326:30/5.
- [12] Bove AA. Introduction. In: Bove AA, editor. Diving Medicine, 3rd ed.. Philadelphia, PA: W B Saunders, 1997:3/11
- [13] Guyton AC, Hall JE. Physiology of deep-sea diving and other hyperbaric conditions, In: Guyton AC, Hall JE, editors. Textbook of medical physiology/Arthur C Guyton, John E. Hall. 10th ed. Philadelphia, PA: Saunders, 2000, pp. 504/509.
- [14] Moon RE. Treatment of diving emergencies. Crit Care Clin 1999;15:429/5
- [15] <u>https://www.epa.gov/pesticide-registration/list-n-disinfectants-use-</u> against-sars-cov-2 [15]Chesire WP, Jr., Ott MC. Headache in divers. Headache2001;41:235/47.



Page 25 of

[16] DAN Europe {https://www.daneurope.org/} COVID19 Preventive Recomendations for diving community.